



Welcome to the CLU-IN Internet Seminar

Bioavailability-Based Remediation of Metals Using Soil Amendments: Considerations & Evaluation Techniques: Part 1

Sponsored by: U.S. EPA Office of Superfund Remediation and Technology Innovation

Delivered: June 22, 2011, 2:00 PM - 4:00 PM, EDT (18:00-20:00 GMT)

Instructors:

*Dr. Rufus Chaney, Senior Research Agronomist, USDA Agricultural Research Service
(Rufus.Chaney@ARS.USDA.GOV)*

Dr. Mark Sprenger, Environmental Scientist, U.S. EPA OSRTI (sprenger.mark@epa.gov)

Michele Mahoney, U.S. EPA OSRTI (mahoney.michele@epa.gov)


Moderator:

*Michele Mahoney, U.S. EPA, Office of Superfund Remediation and Technology Innovation
(mahoney.michele@epa.gov)*

Visit the Clean Up Information Network online at www.cluin.org

Housekeeping

- Please mute your phone lines, Do NOT put this call on hold
 - press *6 to mute #6 to unmute your lines at anytime
- Q&A
- Turn off any pop-up blockers
- Move through slides using # links on left or buttons



- This event is being recorded
- Archives accessed for free <http://clu.in.org/live/archive/>

2

Although I'm sure that some of you have these rules memorized from previous CLU-IN events, let's run through them quickly for our new participants.

Please mute your phone lines during the seminar to minimize disruption and background noise. If you do not have a mute button, press *6 to mute #6 to unmute your lines at anytime. Also, please do NOT put this call on hold as this may bring delightful, but unwanted background music over the lines and interrupt the seminar.

You should note that throughout the seminar, we will ask for your feedback. You do not need to wait for Q&A breaks to ask questions or provide comments. To submit comments/questions and report technical problems, please use the ? Icon at the top of your screen. You can move forward/backward in the slides by using the single arrow buttons (left moves back 1 slide, right moves advances 1 slide). The double arrowed buttons will take you to 1st and last slides respectively. You may also advance to any slide using the numbered links that appear on the left side of your screen. The button with a house icon will take you back to main seminar page which displays our agenda, speaker information, links to the slides and additional resources. Lastly, the button with a computer disc can be used to download and save today's presentation materials.

With that, please move to slide 3.

Introduction to Bioavailability-Based Remediation of Metals Using Soil Amendments: Considerations & Evaluation Techniques: Part 1

Sponsored by OSWER OSRTI TIFSD

Mark Sprenger
OSWER OSRTI TIFSD ERT
sprenger.mark@epa.gov
732-906-6826

There are a large number of sites and large areas where “soils” have lost their functions. These include mining sites, but can include properties which have been impacted through other means.


The end result of the activities or impacts to these properties may be:


- Loss of top soil;
- Loss of soil functions;
- loss of use options.

These may result from:

- Physical loss of soil;
- Loss of soil structure;
- Toxicity.







Because of the scale of the issues surrounding the remediation of these impacted lands there is a need to improve our ability to evaluate the risks which exist.

$$\text{Exposure} \times \text{Hazard} = \text{Risk}$$

The exposure assessment is a critical element of the risk assessment ; how do we evaluate/determine the site specific exposure?

Examples:

Human health risk assessments may do a market basket surveys;

Ecological Risk assessments may conduct field collection of organisms and measures accumulation.

Both are trying to get at location/site specific bioavailability/bioaccessability the actual exposure.

Framework for Metals Risk Assessment



Environmental toxicology has also had a premise that the total concentration of a contaminant is not a good estimator of exposure. The metals framework outlines the issues and highlights that for metals chemical form is critical in determining the toxicity and the exposure.

We need to explore the available techniques for assessing exposure and hazard. We need to make sure we are applying the techniques correctly and interpreting the results correctly. The goal is to remediate the risks effectively; provide protective remedies.

EcoTools

Tools for Ecological Land Reuse

<http://www.cluin.org/ecotools>

EcoTools include:

-  EPA Presentations
-  Principles for Ecological Land Reuse
-  Soil Amendments
-  Terrestrial Carbon Sequestration
-  Plants and Revegetation
-  Growing Gardens in Urban Soils
-  Act Locally
-  Organizations and Resources
-  Land Revitalization Assistance
-  Case Study Profiles



<http://www.cluin.org/ecotools>

Archived Internet Seminars and Presentations

EPA Presentations

Archived internet seminars for ecological restoration

Since 1998, CLU-IN has presented Internet Seminars covering a wide variety of technical topics related to hazardous waste characterization, monitoring, and remediation. For each seminar topic, we have selected the highest-quality offering for placement in our archives.

- NARPM Presents...Ecological Revitalization: Turning Contaminated Properties into Community Assets: Archive of Mar 15, 2011 Seminar
- NARPM Presents...A Tale of Three Sites — Supporting Reuse Throughout the Cleanup Process: Archive of Mar 10, 2011 Seminar
- Superfund Redevelopment Seminar Series: Archive of Sep 30, 2010 Seminar
- Identifying & Evaluating Ecosystem Services at Contaminated Sites Prior to Remediation: Archive of Aug 18, 2010 Seminar
- Green Remediation: Applying Strategies in the Field — Session 1 of 3: Archive of Fall 2009 Seminars
- Ecological Revitalization Resources Available through EPA - Parts 1 and 2: Archive of Dec 3 and 5, 2007 Seminars
- Ecological Revitalization Resources at Various Federal Agencies: Archive of Nov 27, 2007 Seminar
- Ecological Revitalization Case Studies - The Atlas Tack Site and the Poudre River Site: Archive of Aug 2, 2007 Seminar
- Understanding and Reconstructing Soil Conditions at Remediation Sites: Archive of May 2, 2007 Seminar
- Revegetation and Restoration of an Oil Contaminated Wetland in Northern New Jersey: Archive of Dec 14, 2006 Seminar
- Jump-Starting Ecological Restoration - Soil Health: Archive of Oct 5, 2006



EcoTools

Tools for Ecological Land Reuse

<http://www.cluin.org/ecotools>

Fact Sheets



Using Soil Amendments

FACILITATING SITE REMEDIATION, REVITALIZATION, AND REUSE

Superfund sites and other sites with contaminated or disturbed soils exhibit a variety of problems that often can be addressed effectively and directly through the application of soil amendments. The use of soil amendments can be a cost-effective in situ practice for remediation, revitalization, and reuse of many types of disturbed and contaminated landscapes.

When applied and applied properly, soil amendments both reduce the contaminated exposure pathways and reduce soil physicochemistry. Soil amendments also can reduce vegetation soil conditions for plant growth by lowering pH, adding organic matter, restoring soil structure, increasing moisture retention, and reducing erosion. This fact sheet provides an overview of using soil amendments to address on-site contamination and the resources available to assist with this clean-up approach.

DID YOU KNOW?

Technical Assistance for using soil amendments to address contaminated soils is available through the Office of Superfund Remediation and Technology Assistance (OSRTA). For more information, please contact OSRTA's Remedial Technology Branch at (202) 455-5050 or by e-mail at osrta@epa.gov.

Why is using soil amendments effective?

Soil amendments are used to alter the physical and chemical properties of soils. They can be used to reduce the availability of contaminants to plants and animals, to reduce the mobility of contaminants in the soil, and to improve the soil's ability to support vegetation. They can also be used to improve the soil's ability to support vegetation and to improve the soil's ability to support vegetation.

What are soil amendments?

Soil amendments are materials added to soils to improve soil structure and soil fertility. They can be used to improve the soil's ability to support vegetation and to improve the soil's ability to support vegetation.

Notes from the Field

At the Superfund site, the soil was contaminated with lead and arsenic. The soil was amended with compost and lime. The soil was amended with compost and lime. The soil was amended with compost and lime.

Ecological Revitalization Database Fact Sheet

Introduction

The U.S. Environmental Protection Agency (EPA) Office of Superfund Remediation and Technology Assistance (OSRTA), Remedial Technology and Field Services Division (RTFD), is pleased to provide this fact sheet to assist project managers and others with using the Ecological Revitalization Database (ERD) to support the goal of using previously disturbed sites for ecological revitalization. The ERD is a database of sites that have been previously disturbed and are now being used for ecological revitalization. The ERD is a database of sites that have been previously disturbed and are now being used for ecological revitalization.

Background Information

Ecological revitalization sites are the result of a variety of factors, including land use changes, natural processes, and human activities. The ERD is a database of sites that have been previously disturbed and are now being used for ecological revitalization.

How Can I Share Information on Additional Projects and Sites?

ERD is continuing to efforts to increase the use of ecological revitalization. Sites of particular interest include natural resources, agriculture, historic resources, and other sites that are of interest to the public. The ERD is a database of sites that have been previously disturbed and are now being used for ecological revitalization.



REUSING POTENTIALLY CONTAMINATED LANDSCAPES: Growing Gardens in Urban Soils

This fact sheet provides communities and individuals with general information about:

- Common contaminants that can be found in urban soils.
- Ways to identify contaminants and reduce exposure.
- Improving soils and growing plants in urban soils.
- Additional resources and technical assistance.

Introduction

Communities throughout the country are working to reduce exposure and growing in a sustainable space to increase their health, safety, and the quality of life. One of the ways to do this is by growing gardens in urban soils. Growing gardens in urban soils is a way to improve the quality of the soil and to improve the quality of the soil.

More Information for the Urban Gardener

For more information on the urban gardener, visit the website at www.epa.gov/osrta/urban_gardening.

Ecological Revitalization Database



- Site Information
 - Site name, location and cleanup program
 - Entity responsible for cleanup
- Project Information
 - Project name
 - Site history and background
 - Site use prior to revitalization
 - Final use after revitalization – The final use of the land after the revitalization process is complete (open space, ready-for-reuse, wildlife refuge, etc.).
 - Habitats created or restored – Any habitats that are created or restored on an entire property or on a portion of a property through ecological revitalization (wetland, grassland, stream, etc.).
 - Soil amendments – Any soil amendments that are used to aid in soil growth and health. Soil amendments are materials added to soils in order to make them suitable for sustaining plant life or development.
 - Contaminated media type and concentration – The contaminated media as well as the concentration of the contaminant in each specific media.



100+ project profiles!

- Remediation technology – The technology used to remediate the contaminated media and site.
- Remedy description
- Operations and maintenance (O&M) requirements – O&M for the overall cleanup typically includes inspection, sampling and analysis, routine maintenance and small repairs, and reporting, as necessary. Any additional O&M requirements, including those specific to ecological revitalization activities, are listed.
- Long-term stewardship at the site – Long-term stewardship as state voluntary cleanup programs and property owners have primary responsibility for carrying out maintenance of engineering controls and ICs for the long-term.
- Issues faced
- Solutions for issues faced
- Additional Information
 - Points of contact
 - References

Soil Amendments





Site-Specific Support

- Soil Amendments for Remediation & Reuse
- Site Reuse Planning
- Expert consultation
- Documentation
- Presentations
- Other?

EPA Using Soil Amendments
FACILITATING SITE REMEDIATION, REVITALIZATION, AND REUSE
www.epa.gov/soil

Superfund sites and other sites with contaminated or disturbed soils exhibit a variety of problems that often can be addressed effectively and directly through the application of soil amendments. The use of soil amendments can be a cost effective in site process for remediation, revitalization, and reuse of many types of disturbed and contaminated landscapes.

When specified and applied properly, soil amendments limit many of the contaminant exposure pathways and reduce soil phytotoxicity. Soil amendments also can reduce evaporative soil conditions for plant growth by balancing pH, adding organic matter, restoring soil microbial activity, increasing moisture retention, and reducing compaction. This fact sheet provides an overview of using soil amendments to address on-site contamination and the resources available to assist with this clean up approach.

DID YOU KNOW?
Technical Assistance for using soil amendments to address contaminated soils is available through Office of Superfund Remediation and Technology Innovation (OSRTI). For more information, please contact Michele Mahoney by phone at 703-603-9057 or by e-mail at mahoney202@epa.gov.

Why is using soil amendments effective?
Soil amendments can reduce the bioavailability of a wide range of contaminants and, consequently, while amendments affecting co-contaminants occur and, thereby, protecting against future movement of contaminants to soil and water. They can be used to restore long-term site critical contaminant concentrations to long-term ecological restoration projects. Using natural materials (natural hydrophobic) in soil amendments offers the potential for significant cost savings compared to traditional alternatives. In addition, land revitalization using soil amendments has significant ecological benefits including wildlife habitat, species diversity, food control aesthetics, and recreation.

What are soil amendments?
Soil amendments are materials added to soils in order to improve soil quality and establish plant growth. Soil amendments may be inorganic (e.g., lime, organic matter, organic or inorganic) or organic (e.g., lime, organic matter). Commonly used soil amendments include:

- mineral fertilizers, such as water treatment residuals;
- animal manure and compost;
- sewage sludge;
- lime;
- organic matter, such as, or degraded materials;
- log wood chips;
- animal products by products;
- manure and compost;
- composted manure, and
- a variety of commercial agricultural hydrophobic, as well as traditional agricultural fertilizers.

Notes from the Field
At the California State site in Jackson, California, bioremediation along the river channel that has the soil pH and organic concentrations of metals. One way used to amend the soil pH and organic matter content in the bioremediation. A wide range of materials are used, including natural and synthetic materials. Results show that the bioremediation of the river channel is a promising approach to remediate the river channel after being treated with soil amendments and water treatment residuals (WTR).

EPA Environmental Remediation Agency
Office of Superfund Remediation and Technology Innovation



Michele Mahoney

US EPA OSWER

Technology Innovation & Field Services Division

Phone: (703) 603-9057

Email: Mahoney.Michele@epamail.epa.gov

www.cluin.org/ecotools

**Bioavailability-Based Remediation of Soil
Metals Using Soil Amendments:
Considerations & Evaluation Techniques: Part
1**

Rufus L. Chaney

**USDA-Agricultural Research Service
Environmental Management and Byproducts Utilization Lab
Beltsville, MD.**

Webinar, June 22, 2011

The Problem:

- **Highly metal contaminated barren soils**
 - Mine wastes
 - Smelter contamination
 - Contaminated riverine or lake sediments on land
- **Rich in phytotoxic elements**
 - Zn, Cu, Ni, Mn, Co.
- **Risk thru soil ingestion – Pb, As, F**
- **Risk thru food chain transfer – Cd, Mo, Co**
- **May be highly acidic due to oxidation of sulfide**
 - Pyritic mine waste
 - Natural soil acidity coupled with acidic rainfall
- **Need: Reduce bioavailability of soil metal to remediate risks.**



²³
Palmerton, PA, 1980; Dead Ecosystem on Blue Mountain--Zn, Cd, Pb

Bunker Hill, Kellogg, Idaho-Superfund Site – Zn, Cd, Pb



24

Chuck Henry collecting test soil at Leadville, CO site – Zn, Cd, Pb.





**Belvidere Mountain Site, Vermont
Serpentine Asbestos Mine Wastes**



Baltimore Urban Garden, 1980.

Revegetation/Remediation of Heavy Metal Contaminated Soils: Problems.

- **Low soil pH or pH decline from pyrite oxidation.**
 - Make site calcareous; balance Ca and Mg; Mn if needed.
 - Limestone with biodegradable organic matter aids leaching
- **Nutrient Deficiencies, especially P and N.**
 - High Pb soils need higher P addition to precipitate Pb.
 - Higher available soil P needed to maintain legumes.
 - No metal tolerant legumes to supply N to grasses
- **Need more metal sorption by soil, Fe for grasses.**
 - Grasses obtain Fe using secreted phytosiderophores (chelators), so higher soil Fe aids grasses metal resistance
- **Low organic matter, lack of microbes--Zn Toxicity**
 - Biosolids, manures and composts – inexpensive source of Organic Matter and microbial inoculant.

28

Revegetation/Remediation of Heavy Metal Contaminated Soils: Solutions.

- **Make Soil Calcareous Using By-Product Lime**
 - Increases metal adsorption and occlusion.
 - Alleviates phytotoxicity of Zn, Cu, Ni, Cd, etc.
- **Increase Metal Adsorption Capacity**
 - Include Fe, Mn hydrous oxides and phosphate.
 - Provides persistent reduction in metal toxicity.
- **Remediated Soil Must Support Legumes.**
 - High pH and soil P aids legume competition, alleviating need for annual N fertilization.
- **Food Chain Protection: Cd/Zn ratio; calcareous.**
- **Reduced bioavailability of soil Pb, As, Cd, etc. to animals with soil exposure at remediated site.**
- **Effective plant cover reduces soil ingestion.**

29

Tailor-Made Mixtures For Remediation of Metal Toxic Soils

- **Mix Composts, Biosolids and Byproducts to Complement benefits or improve metal sorption:**
 - APL Biosolids and composts
 - Composts of Yard Debris or pre-separated MSW.
 - Agricultural Organic Byproducts
 - Manures; crop residues; food processing byproducts
 - Fe, Mn, Silicate Byproducts from industry.
 - Coal Combustion Byproducts, FGDB, Ash.
 - Drinking Water Treatment Residues
 - Limestone equivalent byproducts.
 - Wood ash; waste lime; sugarbeet lime; fly ash; etc.


30

Tailor-Made Biosolids Mixtures For Beneficial Use and Remediation

- **Apply mixture of limestone equivalent, metal adsorbent, organic soil amendment, and fertilizer value to correct all risks/problems of the contaminated soils:**
 - Zn or Ni Phytotoxicity; make soil calcareous.
 - Food-chain risks from Cd prevented by Zn.
 - Soil ingestion risk from soil Pb, As, etc.
 - N fixation by legumes made possible.
 - Leaching of limestone equivalent corrects surface and subsurface soil metal phytotoxicity.
- **One treatment for comprehensive remediation.**

31

Complex Equilibria of Metal Ions with Components of Soil Environments



SOIL-PLANT BARRIER


Processes in soils or plants which prevent excessive food-chain transfer of elements

- **Insolubility or adsorption in soil or plants roots:**
 - Cr, Pb, Fe, Hg, Sn, Au, Ag, Zr, Al, Ce, Ti, etc.
- **Phytotoxicity limits plant yield at levels which are not toxic for lifetime consumption by livestock:**
 - Zn, Cu, Ni, As, Mn, B, F, etc.
- **Exceptions to Soil-Plant Barrier:**
 - Cd, Se possible risk to humans
 - Mo, Se, Co possible risk to livestock
- **Barrier can be circumvented by direct ingestion of surface soils.**
 - Pb, As, F, Hg, Fe may comprise risk if high on surface.


33

Difficulties in Raising Soil pH


- Agricultural limestone can only react with soil acidity very near the limestone articles.
 - Diffusion of Ca^{2+} and H^+ only short (mm) distances.
- Need to raise pH of contaminated soil depth using mixing or alternatives.
- We found that mixing limestone equivalent with biodegradable organic matter formed alkaline leachable mixture.
 - Ca-organic acid complexes can readily leach.
 - Oxidation of the organic acid essentially leaves a residue of highly reactive CaCO_3 at depth.
 - The greater the rate of biodegradation of the applied organic matter, the greater the leaching of CaCO_3 .
 - Finer lime materials more reactive.



Effect of rates of limed digested biosolids applied to Galestown loamy sand in 1976 on pH at soil depths in 1992 (Brown et al., 1997). 35



Effect of rates of limed digested biosolids applied to Christiana fine sandy loam in 1976 on pH at soil depths in 1992 (Brown et al., 1997).



Effect of limed biosolids or composts applied to **Christiana fine sandy loam** in 1976 on pH at soil depths in 1992 (Brown et al., 1997).

Effect of Biosolids Processing Technology in Reducing Soil-Pb Bioavailability.

- Conducted by Brown, Xue, Hallfrisch and Chaney/WERF.
- Baltimore urban soil = 2135 mg Pb/kg
- Mixed biosolids products at 10% dry weight (224 t/ha) or equivalent added biosolids matrix.
 - Incubated moist for 30 days; dried; mixed.
- Added 5% soil to purified rat diet for 35 day feeding period -- simulates *pica* soil ingestion levels.
 - Measured Pb in blood, bone, kidney, etc.
 - Compare to Pb-acetate (soluble; 100% bioavailable); interpolate Pb-acetate which gives equal tissue Pb concentration as soils.

Effect of Biosolids Processing Technology in Reducing Soil-Pb Bioavailability.

Treatment	Soil Pb	Diet Pb	Bone-Pb
	-----mg/kg dry weight-----		
Unamended	2135	125.3 a	144.6 a
Syracuse Raw	2099	82.4 cd	87.5 b-e
Syracuse Pellet.	2034	84.3 cd	86.7 cde
Syracuse Comp.	1768	116.7 ab	104.5 bc
Orgro-Baltimore	2576	100.6 b	73.3 de
Compro-DC	2309	99.2 b	81.8 cde


**Bioavailability of Cadmium in Biosolids-
Fertilized Swiss Chard Fed at 28% of Diet to
Guinea Pigs for 80 Days (Chaney et al., 1978)**

Treatment	Rate	Soil	Soil	Chard		Kidney	Liver
		Cd	pH	Cd	Zn	Cd	Cd
	t/ha	mg/kg		mg/kg dry		---mg/kg dry---	
Control	0	0.04	6.0	0.5	70	14.9 a	3.1 a
Biosolid-1	56	0.32	5.7	1.5	950	14.5 a	2.7 a
Biosolid-2	112	0.94	5.5	2.7	580	14.5 a	2.7 a
Biosolid-3	224	0.89	6.6	1.4	257	15.8 a	3.6 a



Palmerton, PA, 1980; Dead Ecosystem on Blue Mountain.

41





**View from Hahn farm north of Stoney Ridge
looking toward Blue Mountain-1988.**

43



Foal on Hahn farm showing Zn toxicity, 1979.

44



Proximal tibia showing osteochondrosis, 1979 Foal. 45



Palmerton, PA, 1980; because lawn grasses died from Zn, many residents covered their lawns with stones or mulch.⁴⁶



47

View of West smelter with Blue Mountain in background-1980.



48

Home garden near across railroad from West Smelter, 1980.



49

Chlorotic radishes (Zn phytotoxicity) - Palmerton garden, 1980.



View from Stoney Ridge toward Blue Mountain, 2000. 50



**Palmerton, PA, 1990: Oyler's First Test Plot Using Biosolids +
FlyAsh + Limestone, with 'Merlin' Red Fescue; adjacent control.**

51

**Characteristics of the Blue Mountain North Slope
Soils Sampled in bulk in 1998 for *Thlaspi* studies.**

mg/kg DW

44,100 Zn

25,500 Fe

8,920 Mn

863 Cd

pH 6.25



Palmerton, PA, 1999: Looking down revegetated Blue Mt. 53




Palmerton, PA -- Revegetated Area in 1999: Area with good intermediate wheatgrass and lespedeza cover.

54




**Palmerton, PA: Blue Mountain – 1999; Foreground =
Biosolids+Limestone+FlyAsh; Background = untreated Control**

55



This revegetated landscape is a Superfund cleanup site replanted by Zinc Corporation of America.(K6055-3)



Palmerton, PA 1999; Untreated area adjacent to revegetated area of Blue Mountain, with John Oyler and Tom Stuczynski.

57

Mean total Zn, Cd and Pb, and DTPA-extractable Zn and Cd (at 100 mL extractant/2 g soil) in Palmerton “Revival Field” Test Plots Comparing Traditional and Biosolids Compost Remediation Treatments (Li et al., 2000).

Treatment	Total			DTPA-Extractable	
	Zn	Cd	Pb	Zn	Cd
	----- mg kg ⁻¹ -----				
Control	14900 a†	164. a	687. a	4940. a	83.1 a
Limestone	15700 a	161. a	680. a	4980. a	82.9 a
Compost	16000 a	170. a	767. a	4550. a	69.1 b
†Treatment means followed by the same letter are not significantly different at the 5% level (Duncan-Waller-test).					
Use of DTPA-TEA extraction required using 5 g/50 mL rather than 10 g/20 mL because high soil metals saturated DTPA chelation capacity.					

58


Mean pH, Sr-extractable metals, pH, organic matter and oxalate Extractable Fe and Mn in Palmerton “Revival Field” Plots comparing remediation using traditional or biosolids compost methods; plots Installed in 1993, last sampled in 1998 (Li et al., 2000).

Treatment	<u>Sr(NO₃)₂-Extr.</u>		pH	Organic	<u>Oxalate-Extr.</u>	
	Zn	Cd		Matter	Fe	Mn
	----- mg kg ⁻¹ -----			%	----- g kg ⁻¹ -----	
Control	195. a	1.99 a	5.9	4.6	5.74 a	2.12
Limestone	156. a	1.65 a	6.5	4.7	5.61 a	1.92
Compost	4.8 b	0.033 b	7.2	9.5	16.7 b	2.44
†Treatment means followed by the same letter are not significantly different at the 5% level (Waller-Duncan test.)						




**Revival Field-Palmerton: Yin-Ming Li
and Bev Kershner in ARS photograph.**

60



Palmerton, PA, Revival Field, Year-3: Grasses thrive only on Alkaline Biosolids Compost Treatment (Cooperator Bev Kershner). 61





Appalachian Trail remained barren due to Zn phytotoxicity in 2008.





**Sassafras growing on south face of Blue Mountain near Palmerton, PA., 6-21-2006
Leaves show severe interveinal chlorosis expected from Zn phytotoxicity.**

65

Why Use High Quality Tailor-Made Biosolids Mixtures in Remediation of Soil Metals?

- Fe and phosphate in biosolids increase metal “specific adsorption ability of the soil, reducing metal phytoavailability.”
 - Can remediate Zn phytotoxicity and food chain Cd risk.
 - Can reduce soil Pb bioavailability/form Pb pyromorphite.
- Combining limestone equivalent and biodegradable organic matter causes alkalinity to leach down soil profile.
 - Corrects subsoil acidity and metal phytotoxicity/leachability.
- With pH buffered by applied limestone equivalent, metal adsorption is maximized, and occlusion promoted.
 - Some metals are occluded in crystalline Fe oxides, Mn oxides.
- Organic matter and balanced nutrient supply supports crops!
- Tailor-Made Remediation Mixtures can immediately inactivate metals, provide microbial inoculum, add energy and nutrients.
- Cost savings; public benefit.

66

What Does it Take To Develop Local Tailor-Made Remediation Products?

- Risk assessment and value information from evaluation of field studies of product utilization.
- Courageous agencies and businesspersons who will seek out such combinations of biosolids, byproducts, and valuable commercial uses of the products.
- Organized valid risk assessment information on:
 - Phytoavailability of applied and soil elements in field.
 - Bioavailability of soil and crop elements.
- Improved risk communication, and honest risk assessments. Examples from Cd food-chain risk, soil Pb and As risk, and phytotoxicity risks from biosolids show massive errors of conservative assumptions.

67

Bunker Hill, Kellogg, Idaho-Superfund Site





Bunker Hill, Idaho -- Smelter killed ecosystem Superfund Site.





Aerospreader Applying Biosolids-Wood Ash Mixture at Bunker Hill ⁷⁰



Highly Zn-phytotoxic smelter and mine waste contaminated soils at Bunker Hill, ID (15,000 mg Zn/kg);

Background = Biosolids+Wood-Ash Remediated

Foreground = Seeded control hazardous soil.

71



Revegetation of Bunker Hill Hillsides using mixture of biosolids, woodash and logyard debris, after 2 years.

72

Remediation of Page Swamp

- The Page Swamp is a wetland constructed in a Pb-Zn-Cd mining waste storage pile near Kellogg, ID.
- In cooperation with US-EPA Superfund ERT, Henry and Brown of Univ. Washington, Chaney et al. tested application of organic amendment plus alkaline byproducts to remediate the highly contaminated site soils.
- Before treatment, the site lacked vegetation even when flooded. Further, the acidity allowed soil metals to inhibit soil microbes so that flooded soil did not become sufficiently reducing to form PbS.
- Application of the composted biosolids plus wood ash mixture prevented toxicity to microbes or plants, soil became highly reducing and PbS was formed
 - Formation of PbS reduces risk to birds which ingest sediments.
 - Vegetation was low in metals and safe for wildlife consumption.

73



Page Swamp near Kellogg, ID; barren wetland built in mine wastes;
Mixture of compost and wood ash applied by Aerospreader.



West Page Swamp prior to beginning treatment (10/7/98)



Overview and beginnings of final treatment by blower (9/21/00)



**Page Swamp remediated area in next season after reactions
Of soil amendments and natural plant colonization.**

Design of Experiment to Test Remediation (Phytostabilization) of Ni-Phytotoxic Soil With Limestone and Fertilizers For Crops Which Differ in Susceptibility to Ni.

- Plant species tested

Poaceae

Corn Wheat

Barley Ryegrass

Oat

Dicots

Radish Tomato

Bean Soybean

Redbeet Swiss Chard

- Three limestone rates

0. - Control

2.54 Mg ha⁻¹ - Limed soil (pH 6.0)

50. Mg ha⁻¹ - Calcareous soil (pH 7.7)

Lime=powdered reagent grade $\text{CaCO}_3 + \text{MgCO}_3$ (4.8 : 1 w/w)

77


Experimental Design-2

- **Mn rates – Applied as $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ to prevent Mn deficiency seen on Port Colborne Soils if limed.**
 - Control - 20 kg ha⁻¹
 - Limed soil - 50 kg ha⁻¹
 - Calcareous soil - 100 kg ha⁻¹
- **Plants grown for 42 days .**
- **Ni in plant tissue determined by AAS after ashing and digestion in HNO_3/HCl .**
- **Soil pH measured in 1:2 v/v water slurry-Harvest**
- **Soil Ni extracted with 0.01 M $\text{Sr}(\text{NO}_3)_2$.**
 - 10g soil:40 mL solution (Helmke, Corey et al.);
 - Shaken for 2 hr, filtered; analysis by AAS.

Properties of Welland Clay Loam Soil (from Port Colborne, Ontario, Canada, Used in Nickel Phytotoxicity Remediation Tests.


Measurement	Welland
Total Ni, mg/kg	2900
DTPA-Ni, mg/kg	634
Sr(NO ₃) ₂ -Ni, mg/kg	57
Initial pH	5.2

79



The effect of soil pH on 0.01 M $\text{Sr}(\text{NO}_3)_2$ -extractable soil Ni.

80





82


The Vermont Asbestos Group Site Near Eden/Lowell/Stowe, VT.

- **Has been barren since about 1950. Potential dispersal of asbestos from the barren ground rock presents an environmental risk.**
- **Rock is serpentinite, rich in Ni, Cr, Co, and Mg silicate. Deficient in many plant nutrients.**
- **Extremely infertile; Mg phytotoxic=Ca deficient.**
- **Not Ni, Co or Cr phytotoxic due to high pH (>8.0) which is caused by presence of Mg-silicate.**
- **Alternative to *in situ* phytostabilization would be covering mine waste with 12-24 inches of topsoil!** ⁸³

Severe Infertility and Lack of Soil Properties Prevent Plant Survival

- **Serpentine soils are Mg phytotoxic due to very low Ca:Mg ratio of this type of rock.**
- **N, P, K, and trace elements are also deficient.**
- **Serpentine soils are normally severely Ca and P deficient for all but serpentine ecology plants.**
- **Because site has high slopes, goal was to use surface applied amendment mixture to achieve revegetation at low cost.**
- **Designed experiment to evaluate surface applied compost plus Ca and NPK fertilizers.**

**Belvidere Mountain Site, Vermont
Serpentine Asbestos Mine Wastes**





**Belvidere Mountain Site, Vermont
Serpentine Asbestos Mine Wastes**




**Vermont Asbestos Group Belvidere Mountain Site
Serpentine Asbestos Mine Wastes**

Treatments Tested:


- **Surface Applied Soil Amendments:**
 - Control
 - NPK Fertilizer (normal roadside revegetation)
 - Compost + NPK
 - Compost + NPK + Gypsum(=CaSO₄)
 - Topsoil + NPK
- **Plant Species Tested:**
 - Kentucky bluegrass
 - Perennial ryegrass
 - Tall Fescue
 - Alsike Clover

Tall Fescue

47 Days from Seeding




Perennial Ryegrass 47 Days from Seeding




Perennial Ryegrass

47 Days from Seeding




Kentucky Bluegrass 47 Days from Seeding



Kentucky Bluegrass 47 Days from Seeding



Alsike Clover 47 Days from Seeding



Alsike Clover 47 Days from Seeding





**Preparing mixture of COMPOST (manure and yard debris),
mined gypsum, NPK fertilizer plus limestone**




August 24, 2011: Applying the compost mixtures to test plots; compost was raked even, then seeded with crop mix.



**Test plots with two compost mixtures vs. Control
(three replications in RCB) VAG site August 23, 2010.**

Cover crops establishment -- Sept. 30, 2010 at VAG Site.





Cover crop observed on May 24, 2011

How Did We Achieve Success on VAG Site?

- **Evaluated composition of soil for metals, pH, and nutrients before plant testing.**
- **Recognized severe Ca and P infertility of serpentine rock derived soil materials.**
- **Tested treatments and plant species on site soil in greenhouse.**
- **Amendment mixture included all nutrients needed for plant growth in compost.**
- **Added limestone to prevent acidification of compost layer over time with N-fixation.**
- **Included gypsum to add Ca to sub-surface soil.**

Summary

- **Risk Assessment of contaminated soil:**
 - Soil-Plant Barrier.
 - Phytoavailability related to soluble metal level.
 - Affected by pH, sorbents (Fe, Mn, OM) and competition.
 - Bioavailability of metals in ingested soil requires test correlated with bioavailability to animals.
 - Important risk for Pb, As, F, and some others.
- **In situ remediation using byproducts to reduce phytoavailability, bioavailability and improve agronomy.**
 - Alkalinity to reduce metal solubility.
 - Organic matter/N to improve fertility.
 - Diverse microbial inoculum.
 - **Support growth of perennial grasses and legumes.**

102

What Does it Take To Develop Local Tailor-Made Remediation Products?

- Risk assessment and value information from evaluation of field studies of product utilization.
- Courageous agencies and businesspersons who will seek out such combinations of biosolids, byproducts, and valuable commercial uses of the products.
- Organized valid risk assessment information on:
 - Phytoavailability of applied and soil elements in field.
 - Bioavailability of soil and crop elements.
- Improved risk communication, and honest risk assessments. Examples from Cd food-chain risk, soil Pb and As risk, and phytotoxicity risks from biosolids show massive errors of conservative assumptions.

Summary

- **One Shot Remediation of Metal Toxic Soils:**
 - For Zn, Cu, Ni rich acidic soils causing phytotoxicity.
 - Make contaminated soil depth calcareous
 - Provide enough P, K, and other nutrients to support diverse vegetation, and enough organic-N to achieve stable ecosystem which includes legumes.
 - For Pb or As co-contaminated soils have to reduce bioavailability of Pb or As in ingested soil.
 - Phosphate and composts can reduce soil Pb bioavailability.
 - Iron oxides can reduce soil As bioavailability
 - With normal <1:100 Cd:Zn ratio, Zn limits plant growth before Cd accumulated in plants is a risk to foods.
 - If slope of the site is too high for tillage, can combine biodegradable amendments with alkaline **organic** amendments and surface apply; allow rainfall to leach soluble alkalinity into soil profile.

104

Resources & Feedback

- To view a complete list of resources for this seminar, please visit the [Additional Resources](#)
- Please complete the [Feedback Form](#) to help ensure events like this are offered in the future

The screenshot shows a web form titled "U.S. EPA Technical Support Project Engineering Forum Green Remediation Opening the Door to Field Use Session C (Green Remediation Tools and Examples) Seminar Feedback Form". The form includes fields for "First Name", "Last Name", "Email Address", and "Date of Seminar". A checkbox labeled "Please send a copy of my feedback confirmation as a record of my participation to this address" is highlighted with a red circle. The form also includes a "Delivery Address" field.

Need confirmation of your participation today?

Fill out the feedback form and check box for confirmation email.